

RELATIONSHIP BETWEEN VITAMIN D AND PHYSICAL PERFORMANCE IN COMMUNITY DWELLING ELDERLY MEN

V Środkowo Europejski Kongres Osteoporozy i Osteoartrozy oraz XVII Zjazd Polskiego Towarzystwa Osteoartrologii i Polskiej Fundacji Osteoporozy, Kraków 29.09-1.10.2011

Streszczenia:

Ortopedia Traumatologia Rehabilitacja 2011, vol 13 (Suppl. 1).
str 106

P02

RELATIONSHIP BETWEEN VITAMIN D AND PHYSICAL PERFORMANCE IN COMMUNITY DWELLING ELDERLY MEN

Strazdiene V.², Tamulaitiene M.^{1,3}, Mastaviciute A.^{1,3}, Alekna V.^{1,2}

¹Vilnius University, Faculty of Medicine, Vilnius, Lithuania

²State Research Institute Centre for Innovative Medicine, Vilnius, Lithuania

³National Osteoporosis Center, Vilnius, Lithuania

Keywords: *vitamin D, short physical performance battery, elderly men*

Introduction. Vitamin D deficiency is common in the

elderly population and is associated with poor physical performance. Vitamin D may have attribute to muscle strength through a highly specific nuclear receptor in muscle tissue.

Aim: To investigate the association between vitamin D level and physical performance in older men.

Materials and methods. This was a pilot cross-sectional study on community dwelling men aged 60 years and more who visited National Osteoporosis Center in Vilnius, Lithuania. Serum vitamin D was measured by automated immunoassay (Cobas E411, Roche Diagnostic). The standard threshold was used for vitamin D levels of optimal (≥ 30 ng/ml), insufficient (20-29 ng/ml), deficient (< 20 ng/ml), severe depletion (< 10 ng/ml). Physical performance was assessed by the short physical performance battery (SPPB). The SPPB consists of standing balance tasks, five repeated chair stand test and the 4-m walk test. Each of the three performance components were measured in seconds and a score ranging from 0 to 4 was assigned. The sum of three scores composed the total SPPB score ranging from 0 to 12. Statistical analysis was carried out using SPSS version 18.0 for Windows. Pearson correlation coefficient was used to evaluate the relationship between variables. Significance level was defined as 0.05.

Results. A total of 115 men with a mean age 72.1 ± 6.8 years were investigated. Age was statistically significantly negatively associated with the total score of SPPB (Pearson correlation coefficient = -0.3; $p = 0.02$), and no association was observed between age and vitamin D level. Optimal level of serum 25(OH)D was found in 7 (6.1%) subjects. Thirty four men (29.5%) had vitamin D insufficiency, 47 subjects (40.9%) – deficiency and 27 men (23.5%) had vitamin D severe depletion. Highest SPPB score (10.3 ± 1.6) was in subjects with optimal vitamin D level, and lowest (9.3 ± 1.9) – in severe vitamin D depletion. There was no statistically significant correlation between vitamin D level and total SPPB score. Each of three short physical performance battery components was analyzed

separately. The vitamin D level showed an inverse association with time to complete five repeated chair stand test only in vitamin D deficient subjects ($r = -0.3$; $p = 0.05$).

Conclusions. The serum vitamin D was statistically significantly negatively associated with time to complete five repeated stand test in older men.