GLYCOSAMINOGLYCANS AND AMINOHEXOSES IN SYNOVIAL FLUID OF INJURED KNEE JOINTS

XI Zjazd Polskiego Towarzystwa Osteoartrologii i Polskiej Fundacji Osteoporozy V Krakowskie Sympozjum Osteoporozy Kraków 27-29.09.2001

Streszczenia: Materiały kongresowe: STRESZCZENIA, s178. Druk: Drukarnia Skinder, ISBN — 83-904008-5-5 *oraz* Osteoporosis International 2001; vol. 12 (Suppl 1), s44.

P095

GLYCOSAMINOGLYCANS AND AMINOHEXOSES IN SYNOVIAL FLUID OF INJURED KNEE JOINTS (ZAWARTOŚĆ GLIKOZAMINOGLIKANÓW I HEKSOZOAMIN W PŁYNIE STAWU KOLANOWEGO PO USZKODZENIU WIĘZADŁOWYM)

Krystyna Średzińskal , Janusz Popko 2, Andrzej Gindzieńskil, Krzysztof Zwierz3 Department of General Chemistry1 , Department of Pediatric Orthopaedics2 , Department of Pharmaceutical Biochemistry3 Medical Academy, 15-274 Białystok, J.Waszyngtona 17.

Injuries to the ligaments and menisuc of the knee are associated with increased risk for developing post-traumatic osteoarthritis connected with an increased release of proteoglycans and glycosaminoglycans from the cartilage. There is little information on the proteoglycans and glycosaminoglycans in the synovial fluid of an injured knee joint. The aim of our investigation has been to evaluate the levels of glycosaminoglycans and aminohexoses in knee joint fluids in relation to the time which has passed after injury. Samples of knee joint fluid were collected during routine diagnostic arthroscopy, 3 weeks to 27 months after injury, from 18 patients (13 men and 5 women) aged 17-38 (mean 24.9 years) .In this fluid were determined : proteins, uronic acids and Glycosaminoglycans were released hexosamines. from proteoglycans by papain digestion, and separated on microcolumns with CF-11 in to : hyaluronic acid, heparan sulphate, chondroitin sulphate -4 and -6, and keratan sulphate.

In the synovial fluid from the knee joint we have found the highest levels of hexosamines and glycosaminoglycans occuring at 1 month after trauma, and diminishing after that.