

MULTIMORBIDITY IN WOMEN WITH OSTEOPOROTIC FRACTURES IN THE REPUBLIC OF BELARUS

VI Środkowo Europejski Kongres Osteoporozy i Osteoartrozy oraz
XVII Zjazd Polskiego Towarzystwa Osteoartrologii i Polskiej
Fundacji Osteoporozy, Kraków 25-26.09.2015

P05

MULTIMORBIDITY IN WOMEN WITH OSTEOPOROTIC FRACTURES IN THE REPUBLIC OF BELARUS

Ramanau H., Rudenka E.

Gomel State Medical University, Gomel, Belarus

Belarusian Medical Academy of Post-Graduate Education, Minsk, Belarus

Key words: *low-energy fracture, osteoporosis, multimorbidity*

Introduction. Most people with osteoporosis have multimorbidity, but not of them have fractures. The increasing risk of the fracture could be connecting with some chronic disease.

Aim of the Study. The objective of this study was to

define the odds ratio for low-energy fractures for women aged over 50 with difference chronic diseases.

Materials and Methods. The study protocol included the data about chronic diseases diabetes mellitus (DM), myocardial infarction (MI), any cases of cancers, GI ulcers, stroke, COPD, heart failure (HF) and rheumatoid arthritis (RA). The low-energy fractures of forearm, hip fractures and spine was taking account for all female patients aged over 50.

Results. During the research study it was revealed 416 previous low-energy fractures of typical localization among 1533 women with osteoporosis. According to the results of the statistical analysis in patients with fractures of the most frequent combination of osteoporosis and COPD (OR=5.98; 95% CI 2.72-12.60; $p<0.001$), RA (OR=4.11; 95% CI 2.53-6.64; $p<0.001$), HF (OR=2.47; 95% CI 1.70-3.58; $p<0.001$), stroke (OR=2.40; 95% CI 1.46-3.95; $p<0.001$), DM (OR=1.64; 95% CI 1.14-2.39; $p<0.05$) and GI ulcers (OR=1.46; 95% CI 1.07-2.01; $p<0.05$). There are now significant differences between groups with or without fractures if patients had a previous history of MI (OR=1.27; 95% CI 0.68-2.46; $p=0.594$) and any localization of cancer (OR=1.08; 95% CI 0.71-1.67; $p=0.818$).

Conclusions. The presence of some chronically disease in osteoporotic patients may increase the risk of subsequent low-energy fractures in future. These results should be considered for planning strategies antiosteoporotic therapy.

P05

**WIELOCHOROBOWOŚĆ U BIAŁORUSKICH KOBIET ZE ZŁAMANIAM I
OSTEOPOROTYCZNYMI**

Ramanau H., Rudenka E.

Gomel State Medical University, Gomel, Belarus

Belarusian Medical Academy of Post-Graduate Education, Minsk, Belarus

Słowa kluczowe: złamanieiskoenergetyczne, osteoporoza,
wielochorobowość